

83222-22

2/28/2013

1/16



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Direct AG Source, LLC
c/o Jane Miller
Biologic, Inc.
115 Obtuse Hill Road
Brookfield, CT 06804

FEB 28 2013

Product Name: Mepiquat Chloride 4.2%
EPA Reg. No.: 83222-22
Subject: Notification dated February 6, 2013
EPA Decision Number: 475329

Dear Ms. Miller:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10.

The Agency acknowledges the revised labeling to include Kansas in the Directions for Use, as well as the updated emergency contact information and other minor revisions.

The label submitted with the application has been stamped "Notification" and will be placed in our records. If you have any questions, please contact Erin Malone at (703) 347-0253 or via email at malone.erin@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Dris Benmhand".

Dris Benmhand
Acting Product Manager (20)
Fungicide Branch
Registration Division (7504P)



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 83222-22	2. EPA Product Manager T. Kish	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Mepiquat Chloride 4.2%	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Direct AG Source, LLC c/o Biologic, Inc. 115 Obtuse Hill Road Brookfield, CT 06804 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

This is a notification to: 1) add the state of Kansas to the label;; 2) up-date the emergency telephone numbers in the First Aid box; 3) make minor text corrections per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and the EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted				<input type="checkbox"/> Other (Specify) _____	
	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> on label	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		
			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Jane M. Miller	Title Agent	Telephone No. (Include Area Code) (203) 740-1200
------------------------	----------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received

(Stamped)

2. Signature 	3. Title Agent	6. Date Application Received (Stamped)
4. Typed Name Jane M. Miller	5. Date February 6, 2013	

3/16

BIOLOGIC CONSULTING, INC.
agribusiness professionals

2/12

February 6, 2013

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
US Environmental Protection Agency
One Potomac Yard
2777 S. Crystal Drive
Room S-4900, 4th Floor
Arlington, VA 22202

Attention: Mr. T. Kish (PM#22)

Re: "Mepiquat Chloride 4.2%"; EPA Reg. No. 83222-22
Notification to add a state, update emergency contact information, and make other text corrections

Dear Mr. Kish:

On behalf of Direct AG Source please find enclosed this Application for Pesticide Notification to add the following:

1. Add the state of Kansas to appear within the use directions on pages 4, 9, and 10. This state appears on the Me-Too labels for "Mepiquat Chloride 4.2% Liquid", EPA Reg. No. 66222-133 and "Mepichlor 4.2% Liquid", EPA Reg. No. 66330-241.;
2. Update the emergency phone numbers in the "First Aid" section on page 2; and
3. Make minor text corrections on page 1, 4, 5, 9 and 12

This action fits the definition of a Notification and:


"This notification is consistent with the provisions of PR Notice 98-10 and the EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

The following documents are enclosed to process this Notification:

1. Application for Pesticide Notification (8570-1)
2. One (1) copy of "Mepiquat Chloride 4.2% Liquid", EPA Reg. No. 66222-133 and "Mepichlor 4.2% Liquid", EPA Reg. No. 66330-241;
3. One (1) paper copy of the label marked with the added text underlined and highlighted, and the deleted text lined out.
4. One (1) CD of the clean labeling in Pdf format

Should you have any questions, or wish to reach me, please feel free to contact our office at 203-740-1200.

Sincerely,



Jane M. Miller
Agent to Direct AG Source, LLC

```

c c c c c
c c c c c
c c c c c
c c c c c
c c c c c
c c c c c
c c c c c
c c c c c
c c c c c
c c c c c

```

4/16

(FRONT BOTTLE LABEL)

MEPIQUAT CHLORIDE 4.2%

[alt. brand name of FLAT TOP MC]
Plant Growth Regulator

NOTIFICATION

FEB 28 2013

For Use On Cotton

Active Ingredient:

Mepiquat Chloride: N,N-dimethylpiperidinium chloride.....4.2%

Other Ingredients:.....95.8%

Total.....100.0%

*Equivalent to 0.35 pounds of active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside booklet for complete **Precautionary Statements** and **Directions for Use**.


EPA Registration Number 83222-22

EPA Establishment Number

NET CONTENTS:

LOT NUMBER:

Manufactured For:
Direct Ag Source, LLC
30473 260th Street
Eldora, IA 50627



(BOOKLET)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

FIRST AID	
If in eyes:	Hold eyelid open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Hot Line Number	
Have product container or label with you when calling a poison control center or doctor or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday –Friday, 7:30 AM – 3:30 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night for emergency medical treatment information.	

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as nitrile, butyl, neoprene and/or barrier laminate)
- Shoes plus socks

User Safety Requirements

Follow the manufacturer’s instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticide [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.


Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls and long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks.



Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Do not store below 32° F or above 100° F. Store in a dry place away from heat or open flame.

Pesticide Disposal: Pesticide wastes are toxic. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide spray mix, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation of the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable Container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer for recycling, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.


Containers 5 gallons or less: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

I. PRODUCT INFORMATION

MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] is a foliar applied plant regulator that modifies the cotton plant in several beneficial ways. It allows the grower to manage the cotton plant for **short-season production** leading to reduced risk of yield and quality loss due to delayed and prolonged harvest. Additional benefits derived from the use of this product include:

- height reduction and more canopy
- better early boll retention and/or larger bolls
- less boll rot
- improved defoliation
- reduced trash and lower ginning costs
- better harvest efficiency
- darker green leaf color

These benefits can provide for earlier maturity and may result in improved yields.



Spray Coverage

Under most circumstances, water is the recommended diluent, however, oil is permitted in the following states for ultra low volume (ULV) aerial applications: Alabama, Arkansas, Florida, Georgia, Kansas, Louisiana, Missouri, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, and Texas. Refer to **Air and Ground Application** sections for spray volumes.

Regardless of method or gallonage of application, thorough coverage of the cotton foliage is required.

Cleaning Application Equipment


Clean application equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product, particularly if a product with the potential to injure crops was used.

II. APPLICATION INSTRUCTIONS

Early Application

On both short-staple and Pima cotton, the grower has the option of low-rate multiple applications (see **Table 1**) or higher, less frequent applications (see **Table 2**) which greatly facilitates his management flexibility. The multiple application option gives the producer the ability to discontinue usage of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** if any significant stresses occur after an earlier application. In such a case, the total quantity of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** used over a season may be reduced. If stress is relieved, the grower has the option of continuing treatments with **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]**. In addition, the rate and timing ranges indicated in the **Application Rates and Timings Tables** allow the grower to tailor his usage of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** to the degree of vegetative vigor in a given field. In areas where insecticides, miticides or foliar fertilizers are frequently applied, the timings are such that tank mixing is often possible. (See section **VII. Restrictions and Limitations**).

Fields should be carefully scouted and **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** should not be applied if plants are under severe stress from weather factors, mite, insect or nematode damage, disease stress, herbicide injury, or fertility stress. In the absence of these stresses, up to 5 low-rate multiple applications can be made each season. After the first application (at matchhead square in the absence of stress), the rate and timing of subsequent applications will depend on vegetative vigor. Under good growing conditions, additional treatments should be made at 7-14 day intervals. However, if new growth at any time is excessive, higher rates of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** can be used. If significant loss of squares or young bolls has occurred earlier due to insect pressure or other stresses, but now these stresses have all been alleviated, the need for **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** is increased – excess vegetative growth is likely because of poor boll load.



Late Season Application

Late application of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** (approximately during the fourth to sixth week of blooming) can provide certain benefits to cotton. However, it should not and does not substitute for early season use – the time of the greatest benefit from the use of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]**. Late season application can lead to one or more of the following:

- reduction in late season vegetative growth or regrowth after cutout or defoliation
- more complete and manageable cutout
- better defoliation
- earlier maturity
- reduction in trash
- lower ginning costs

Some of these effects may favorably influence the yield potential and fiber quality. A late season application of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** should be applied only if fields are not drought or nutrient stressed; that is, those fields likely to experience additional vegetative growth or regrowth. However, fields that are very rank and extremely vigorous due to a combination of poor boll load and excellent growing conditions may not respond as much as desired to late season applications at the specified rates.

Timing for Late Season Applications

- **On fields where cotton cuts out and then starts regrowth:** Apply when regrowth begins, as evidenced by new leaves in the terminal and stem elongation. This application time is often, but not always, 5-6 weeks after the first bloom.
- **On fields where cotton never completely cuts out:** Apply **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** when there are 4-6 nodes above the white flower (NAWF). Measure NAWF by counting the number of mainstem nodes from the first position white bloom (the one closest to the mainstem) to the terminal. Count the node with the first position white bloom as zero and the last node in the terminal, which is counted, should have a leaf at least the size of a quarter. Generally, the NAWF first reaches 4-6 nodes during the fourth to sixth week of bloom.

During this time, the NAWF should be decreasing about one node every 5-6 days – if its rate of decrease is less, the plant is not cutting out soon enough (the crop is too vigorous). If the fifth week of bloom arrives and NAWF is still above 5-6, apply **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]**.

Use Rate for Late Season Application

Apply 8-24 fluid ounces of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** per acre. Use the lower rate on cotton with only moderate additional growth potential, and the higher rate on fields likely to continue vigorous growth.

AIR APPLICATION

Spray Volume

- **Water as Diluent:** Use a minimum of 2 gallons of water per acre in all states except California. In California, use a minimum of 5 gallons of water per acre.
- **Oil as Diluent:** Use a minimum of 1 quart of oil per acre. When using oil as a diluent, the oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:
 - be nonphytotoxic
 - contain only EPA-exempt ingredients
 - provide good mixing quality in the jar test
 - be successful in local experience

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. If the oil does not contain an emulsifier, one must be added during mixing at a volume equal to 3% of the final volume of the mixing tank. Do not apply **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]**

by ULV without using emulsifiers. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For additional information, see **Compatibility Test for Mix Components**.

AERIAL APPLICATION METHODS AND EQUIPMENT

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the publication titled A Summary of Aerial Application Studies by the Spray Drift Task Force.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying the larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (**see Wind, Temperature and Humidity, and Temperature Inversion section of this label**).

Controlling Droplet Size

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Use a minimum of 2 gallons of water per acre. Increase water volume to at least 10 gallons of water per acre if grass foliage or crop canopy is dense.

Pressure – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure. Use up to 40 psi.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released backward, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types. Use only diaphragm-type nozzles that produce fan spray patterns.

11/16

Boom Length – For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application – Applications should not be made at a height greater than 10 feet above the top of the largest plants. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Do not apply **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** by aircraft when wind is blowing more than 10 mph. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set equipment up to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light and variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent areas, (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g. when wind is blowing away from the sensitive areas). Do not apply **MEPIQUAT PGR** by air if sensitive species are within 200 feet.

GROUND APPLICATION

Spray Volume

- **Water as Diluent:** Use 2 gallons of spray solution per acre in all states except California. In California, use a minimum of 5 gallons of spray solution per acre.

Table 1. Application Rates and Timings: Low Rate Multiple Applications

The times and rates of application have been carefully researched and the **Directions for Use** must be observed as specified below. See section **VI. Restrictions and Limitations**.

Geographic Area	Time of Application	Fields with Moderate Vegetative Vigor: Rate Per Acre	Fields with High Vegetative Vigor: Rate Per Acre
AL, AR, AZ, CA, FL, GA, KS, LA, MO, MS, NC, NM, OK, SC, TN, TX, VA	First Application: Optimal results will be achieved when plants are in the matchhead square ¹ stage of growth	2 fluid ounces	4 fluid ounces
	Second Application: 7-14 days later, or when regrowth occurs.	2 fluid ounces	4 fluid ounces
	Third Application: 7-14 days later, or when regrowth occurs.	2-4 fluid ounces ²	4-8 fluid ounces ²
	Fourth Application: 7-14 days later, or when regrowth occurs.	2-8 fluid ounces ²	4-12 fluid ounces ²
	Fifth Application (if needed): 7-14 days later, or when regrowth occurs.	4-8 fluid ounces ²	4-12 fluid ounces ²
	Late Season: Refer to Late Season Application in section II. Application Instructions .	8-16 fluid ounces ²	12-24 fluid ounces ²
¹ Matchhead square is when the first square of a typical cotton plant is 1/8-1/4 inches in diameter. The first application should be applied when 50% of the plants have one or more matchhead squares. ² Use higher rates if previous application was not made or if growing conditions are conducive to vigorous growth.			




Table 2. Application Rates and Timing: High Rate, Less Frequent Applications

The times and rates of application have been carefully researched and section II. **Application Instruction** must be observed as specified below. See section VI. **Restrictions and Limitations.**

Geographic Area	Time of Application	Rate Per Acre
AL, AR, AZ, CA, FL, GA, LA, MO, MS, NM, NC, SC, TN, VA	<p>First Application: Apply MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] to actively growing cotton that is 20-30" tall, provided cotton is not more than 7 days beyond early bloom stage (5-6 blooms per 25 row feet). If cotton is 24" tall and has no blooms, apply MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]. Use 8-16 fluid ounces per acre on cotton where excessive vegetative growth is not likely to be a problem, and 16 fluid ounces per acre in areas tending to have excessive vegetative growth.</p> <hr/> <p>Second Application for Control of Excessive Vegetative Growth: If the cotton field has a history of vigorous growth or if conditions after the first application of MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] favor vigorous growth, make a second application 2-3 weeks after the first application.</p> <hr/> <p>Third Application for Control of Excessive Vegetative Growth: If the cotton field has a history of vigorous growth or if conditions continue to favor vigorous growth, make a third application 1-2 weeks after the second application.</p> <hr/> <p>Late Season Application: Refer to Late Season Application in section II. Application Instructions.</p>	<p>8-16 fluid ounces</p> <hr/> <p>8-16 fluid ounces</p> <hr/> <p>8-16 fluid ounces</p> <hr/> <p>8-24 fluid ounces</p>
KS, OK, TX (except Rio Grande Valley)	<p>Areas Where Excessive Vegetative Growth is Not a Problem</p> <p>First Application: Apply MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] to actively growing cotton in the early bloom stage (5-6 blooms per 25 row feet). If no blooms are present and the cotton is 20" tall and actively growing, apply MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC].</p> <hr/> <p>Second Application: If conditions after the first application of MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] favor vigorous growth, make a second application 2-3 weeks after the first application.</p> <hr/> <p>Third Application: If conditions after the second application of MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] continue to favor vigorous growth, make a third application 1-2 weeks after the second application.</p> <hr/> <p>Late Season Application: Refer to Late Season Application in section II. Application Instructions.</p>	<p>8 fluid ounces</p> <hr/> <p>8 fluid ounces</p> <hr/> <p>8 fluid ounces</p> <hr/> <p>8-24 fluid ounces</p>
KS, OK, TX (including Rio Grande Valley)	<p>Areas Where Excessive Vegetative Growth is a Problem</p> <p>First Application: Apply MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] to actively growing cotton that is 20-30" tall, provided cotton is not more than 7 days beyond early bloom stage (5-6 blooms per 25 row feet). If cotton is 24" tall and has no blooms, apply MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC].</p> <hr/> <p>Second Application for Control of Excessive Vegetative Growth: If cotton field has a history of vigorous growth, or conditions after the first application of MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] favor vigorous growth, make a second application 2-3 weeks after the first application.</p>	<p>16 fluid ounces</p> <hr/> <p>8-16 fluid ounces</p>

	Third Application: If conditions after the second application of MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] continue to favor vigorous growth, make a third application 1-2 weeks after the second application.	8-16 fluid ounces
	Late Season Application: Refer to Late Season Application in section II. Application Instructions.	8-24 fluid ounces


III. ADDITIVES

If rain is expected within 8 hours, use a high-quality EPA-exempt surfactant to make **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** rain-safe after 4 hours.

Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

1. **Water** – for 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. **Products in PVA Bags** – Cap the jar and invert 10 cycles.
3. **Water-Dispersible Products** – (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) Cap the jar and invert 10 cycles.
4. **Water-Soluble Products** – (such as **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]**) Cap the jar and invert 10 cycles.
5. **Emulsifiable Concentrates** – (oil concentrate) Cap the jar and invert 10 cycles.
6. **Water-Soluble Additives** – Cap the jar and invert 10 cycles.
7. Let the solution stand for 15 minutes.
8. **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, not thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.



IV. MIXING ORDER

1. **Water:** Begin by agitating a thoroughly clean sprayer tank half full of clean water.
2. **Products in PVA Bags:** Rinse the tank thoroughly before adding any material in PVA bags as boron residue will prevent adequate mixing. Place the water-soluble PVA bag into the mixing tank. The water-soluble PVA bag will dissolve in water to allow the contents to disperse. Wait until all water-soluble PVA bags have fully dissolved and the plant regulator is evenly mixed in the spray tank before continuing.
To prepare spray solution for aerial application, use a mixing tank or mixing vat first to get the product into suspension before transferring suspension to air application equipment.
3. **Water-Dispersible Products:** (dry flowables, wettable-powders, suspension concentrates, or suspo-emulsions).
4. **Water-Soluble Products**
5. **Emulsifiable Concentrates**
6. Remaining quantity water.

Only moderate agitation should be used while mixing and transporting.

V. TANK MIXING INFORMATION

MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] has an aqueous base, and as such, is compatible with most insecticides and miticides. You may combine **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** with foliar fertilizers if prior experience has shown the original liquid formulation of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** to be compatible and noninjurious under your conditions. Always perform a **Compatibility Test for Mix Components** before preparing a tank mix application.

Read and follow the applicable **Restrictions and Limitations** and **Directions for Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

VI. RESTRICTIONS AND LIMITATIONS

- Maximum seasonal use rate: Do not apply more than a **total of 48 fluid ounces (3 pints) of MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC] plant regulator** (0.132 pounds a.i.) per acre, per season.
- The sum of all products and formulations containing mepiquat chloride must not exceed **0.132 pounds** of mepiquat chloride per acre per season.
- **Preharvest Interval (PHI):** Do not apply within **30 days** of harvest.
- **Restricted Entry Interval (REI): 12 hours**
- Do not plant another crop within 75 days of last treatment.
- **Stress:** Do not apply to cotton plants under severe stress due to adverse weather conditions, mite, insect, or nematode damage, disease, herbicide injury, or fertility stress. If using the low-rate multiple option, discontinue use until the stress is alleviated. Do not apply a single application of 8-16 fluid ounces of **MEPIQUAT CHLORIDE 4.2% [FLAT TOP MC]** to cotton that is stressed due to lack of soil moisture.
- Do not graze or feed cotton forage to livestock.
- Do not apply through any type of irrigation equipment.




Table 3. Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Cotton	30 days	24 fluid ounces (1.5 pints)	48 fluid ounces (3 pints)	No	Yes

**IMPORTANT INFORMATION - READ BEFORE USING PRODUCT
LIMITATION OF WARRANTY AND LIABILITY**

Read the entire Directions for Use, Conditions of Warranty and Limitations of Liability before using this product. If the terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. - However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Direct AG Source, LLC or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of Direct AG Source, LLC. To the extent consistent with applicable law, all such risks shall be assumed by Buyer or User.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DIRECT AG SOURCE, LLC MAKES NO WARRANTIES, EXPRESS OF IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Direct AG Source, LLC, is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Direct AG Source, LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT DIRECT AG SOURCE, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

